



# Inflammation following acute myocardial infarction: Multiple players, dynamic roles, and novel therapeutic opportunities

Sang-Bing Ong<sup>a,b</sup>, Sauri Hernández-Reséndiz<sup>a,b</sup>, Gustavo E. Crespo-Avilan<sup>a,b</sup>, Regina T. Mukhametshina<sup>g,h</sup>, Xiu-Yi Kwek<sup>a,b</sup>, Hector A. Cabrera-Fuentes<sup>a,b,e,f,g,1</sup>, Derek J. Hausenloy<sup>a,b,c,d,i,j,\*,1</sup>

<sup>a</sup> Cardiovascular and Metabolic Disorders Program, Duke-National University of Singapore, Singapore

<sup>b</sup> National Heart Research Institute Singapore, National Heart Centre Singapore, Singapore

<sup>c</sup> The Hatter Cardiovascular Institute, Institute of Cardiovascular Science, University College London, UK

<sup>d</sup> The National Institute of Health Research University College London Hospitals Biomedical Research Centre, UK

<sup>e</sup> Institute of Biochemistry, Medical School, Justus-Liebig University, Giessen, Germany

<sup>f</sup> Escuela de Ingeniería y Ciencias, Centro de Biotecnología-FEMSA, Tecnológico de Monterrey, Monterrey, NL, Mexico

<sup>g</sup> Institute of Fundamental Medicine and Biology, Kazan (Volga Region) Federal University, Kazan, Russian Federation

<sup>h</sup> German Center for Lung Research, Excellence Cluster Cardio-Pulmonary System, Universities of Giessen and Marburg Lung Center, Giessen, Hessen, Germany

<sup>i</sup> Yong Loo Lin School of Medicine, National University Singapore, Singapore

<sup>j</sup> Barts Heart Centre, St Bartholomew's Hospital, London, UK

## ARTICLE INFO

Available online 9 January 2018

### Keywords:

Inflammation  
Acute myocardial ischemia and reperfusion injury  
Acute myocardial infarction  
Monocytes  
Macrophages  
Lymphocytes  
Dendritic cells  
Cytokines  
Chemokines  
Innate immunity

## ABSTRACT

Acute myocardial infarction (AMI) and the heart failure that often follows, are major causes of death and disability worldwide. As such, new therapies are required to limit myocardial infarct (MI) size, prevent adverse left ventricular (LV) remodeling, and reduce the onset of heart failure following AMI. The inflammatory response to AMI, plays a critical role in determining MI size, and a persistent pro-inflammatory reaction can contribute to adverse post-MI LV remodeling, making inflammation an important therapeutic target for improving outcomes following AMI. In this article, we provide an overview of the multiple players (and their dynamic roles) involved in the complex inflammatory response to AMI and subsequent LV remodeling, and highlight future opportunities for targeting inflammation as a therapeutic strategy for limiting MI size, preventing adverse LV remodeling, and reducing heart failure in AMI patients.

© 2018 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

## Contents

1. Introduction	74
2. The inflammatory response to acute myocardial IRI	74
3. Therapeutic targeting of inflammation following MI	80
4. Conclusion	83
Conflict of interest	83
Acknowledgments	83
References	83

**Abbreviations:** ACS, Acute coronary syndrome; AMI, Acute myocardial infarction; AGEs, Advanced glycation end-products; BAFF, B-cell activating factor; C1-INH, C1-inhibitor; CCL2, Chemokine ligand 2; CCL5, Chemokine ligand 5; CCR2, Chemokine receptor 2; CCR5, Chemokine receptor 5; CCR9, Chemokine receptor 5; CR1, Complement receptor 1; CINC-1, CXCL1, GRO  $\alpha$ , KC, Cytokine-induced neutrophil chemoattractant 1; DAMPs, Damage-associated molecular patterns; ECM, Extracellular matrix; EDA, Extra domain A; eRNA, Extracellular ribonucleic acids; FN-EDA, Fibronectin-end domain A; HSPs, Heat shock proteins; hs-CRP, High-sensitivity C-reactive protein; HMGB1, High mobility group box 1; ICAM-1/ICAM-2, Intercellular adhesion molecule; IFN- $\gamma$ , Interferon- $\gamma$ ; IRF5, Interferon regulatory factor 5; IHD, Ischemic heart disease; IL-1, Interleukin-1; IL-8, CXCL8, Interleukin-8; LV, Left ventricular; LTB4, Leukotriene B4; MIP-2 $\alpha$ , CXCL2, GRO  $\beta$ , Macrophage inflammatory protein-2 $\alpha$ ; MMPs, Matrix metalloproteinases; MCP-1, Monocyte chemoattractant protein-1; MyD, Myeloid differentiation primary response gene; MI, Myocardial infarction; NO, Nitric oxide; NLRs, NOD-like receptors; NF- $\kappa$ B, Nuclear factor kappa-light-chain-enhancer of activated B cells; NLRP3, Nucleotide-binding oligomerization domain-like receptor family of cytosolic proteins; PRRs, Pattern recognition receptors; PS, Phosphatidylserine; PMN, Polymorphonuclear leukocytes; PPCLI, Primary percutaneous coronary intervention; ROS, Reactive oxygen species; RAGE, Receptor for advanced glycation end-products; Tregs, Regulatory T cells; STEMI, ST segment elevation myocardial infarction; TLRs, Toll-like receptors; TGF- $\beta$ , Transforming growth factor- $\beta$ ; TNF $\alpha$ , Tumor necrosis factor- $\alpha$ .

\* Corresponding author at: Cardiovascular & Metabolic Diseases Program, Duke-National University of Singapore, 8 College Road, 169857, Singapore.

E-mail address: [derek.hausenloy@duke-nus.edu.sg](mailto:derek.hausenloy@duke-nus.edu.sg) (D.J. Hausenloy).

<sup>1</sup> These two authors are joint senior authors.